

- mellitus and echocardiographic left ventricular function in free-living elderly men and women: The Cardiovascular Health Study. *Am Heart J*, 133(1), 36-43. doi: 10.1016/s0002-8703(97)70245-x
- 35- Lee, S. J., Kang, J. G., Ryu, O. H., Kim, C. S., Ihm, S. H., Choi, M. G., . . . Kim, T. W. (2009). Effects of alpha-lipoic acid on transforming growth factor beta1-p38 mitogen-activated protein kinase-fibronectin pathway in diabetic nephropathy. *Metabolism*, 58(5), 616-623. doi: 10.1016/j.metabol.2008.12.006
- 36- Lee, T. I., Kao, Y. H., Chen, Y. C., Huang, J. H., Hsiao, F. C., & Chen, Y. J. (2013). Peroxisome proliferator-activated receptors modulate cardiac dysfunction in diabetic cardiomyopathy. *Diabetes Res Clin Pract*, 100(3), 330-339. doi: 10.1016/j.diabres.2013.01.008
- 37- Lee, T. W., Bai, K. J., Lee, T. I., Chao, T. F., Kao, Y. H., & Chen, Y. J. (2017). PPARs modulate cardiac metabolism and mitochondrial function in diabetes. *J Biomed Sci*, 24(1), 5. doi: 10.1186/s12929-016-0309-5
- 38- Lehrke, M., & Marx, N. (2017). Diabetes Mellitus and Heart Failure. *Am J Cardiol*, 120(1S), S37-S47. doi: 10.1016/j.amjcard.2017.05.014
- 39- Lei, S., Li, H., Xu, J., Liu, Y., Gao, X., Wang, J., . . . Xia, Z. (2013). Hyperglycemia-induced protein kinase C beta2 activation induces diastolic cardiac dysfunction in diabetic rats by impairing caveolin-3 expression and Akt/eNOS signaling. *Diabetes*, 62(7), 2318-2328. doi: 10.2337/db12-1391
- 40- Leone, T. C., Weinheimer, C. J., & Kelly, D. P. (1999). A critical role for the peroxisome proliferator-activated receptor alpha (PPARalpha) in the cellular fasting response: the PPARalpha-null mouse as a model of fatty acid oxidation disorders. *Proc Natl Acad Sci U S A*, 96(13), 7473-7478. doi: 10.1073/pnas.96.13.7473
- 41- Li, Z., Abdullah, C. S., & Jin, Z. Q. (2014). Inhibition of PKC-theta preserves cardiac function and reduces fibrosis in streptozotocin-induced diabetic cardiomyopathy. *Br J Pharmacol*, 171(11), 2913-2924. doi: 10.1111/bph.12621
- 42- Lind, M., Bounias, I., Olsson, M., Gudbjornsdottir, S., Svensson, A. M., & Rosengren, A. (2011). Glycaemic control and incidence of heart failure in 20,985 patients with type 1 diabetes: an observational study. *Lancet*, 378(9786), 140-146. doi: 10.1016/S0140-6736(11)60471-6
- 43- Liu, F., Song, R., Feng, Y., Guo, J., Chen, Y., Zhang, Y., . . . Xiao, R. P. (2015). Upregulation of MG53 induces diabetic cardiomyopathy through transcriptional activation of peroxisome proliferation-activated receptor alpha. *Circulation*, 131(9), 795-804. doi: 10.1161/CIRCULATIONAHA.114.012285
- 44- Liu, W., Chen, P., Deng, J., Lv, J., & Liu, J. (2017). Resveratrol and polydatin as modulators of Ca(2+)

- mobilization in the cardiovascular system. *Ann N Y Acad Sci*, 1403(1), 82-91. doi: 10.1111/nyas.13386
- 45- Lytvyn, Y., Bjornstad, P., Udell, J. A., Lovshin, J. A., & Cherney, D. Z. I. (2017). Sodium Glucose Cotransporter-2 Inhibition in Heart Failure: Potential Mechanisms, Clinical Applications, and Summary of Clinical Trials. *Circulation*, 136(17), 1643-1658. doi: 10.1161/CIRCULATIONAHA.117.030012
- 46- Malhotra, A., & Sanghi, V. (1997). Regulation of contractile proteins in diabetic heart. *Cardiovasc Res*, 34(1), 34-40. doi: 10.1016/s0008-6363(97)00059-x
- 47- Mariappan, N., Elks, C. M., Sriramula, S., Guggilam, A., Liu, Z., Borkhsenius, O., & Francis, J. (2010). NF-kappaB-induced oxidative stress contributes to mitochondrial and cardiac dysfunction in type II diabetes. *Cardiovasc Res*, 85(3), 473-483. doi: 10.1093/cvr/cvp305
- 48- Marso, S. P., Bain, S. C., Consoli, A., Eliaschewitz, F. G., Jodar, E., Leiter, L. A., . . . Investigators, S.-. (2016). Semaglutide and Cardiovascular Outcomes in Patients with Type 2 Diabetes. *N Engl J Med*, 375(19), 1834-1844. doi: 10.1056/NEJMoa1607141
- 49- Marso, S. P., Daniels, G. H., Brown-Frandsen, K., Kristensen, P., Mann, J. F., Nauck, M. A., . . . Investigators, L. T. (2016). Liraglutide and Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med*, 375(4), 311-322. doi: 10.1056/NEJMoa1603827
- 50- Miller, J. A. (1999). Impact of hyperglycemia on the renin angiotensin system in early human type 1 diabetes mellitus. *J Am Soc Nephrol*, 10(8), 1778-1785.
- 51- Miller, J. A., Floras, J. S., Zinman, B., Skorecki, K. L., & Logan, A. G. (1996). Effect of hyperglycaemia on arterial pressure, plasma renin activity and renal function in early diabetes. *Clin Sci (Lond)*, 90(3), 189-195. doi: 10.1042/cs0900189
- 52- Murdoch, C. E., Chaubey, S., Zeng, L., Yu, B., Ivetic, A., Walker, S. J., . . . Shah, A. M. (2014). Endothelial NADPH oxidase-2 promotes interstitial cardiac fibrosis and diastolic dysfunction through proinflammatory effects and endothelial-mesenchymal transition. *J Am Coll Cardiol*, 63(24), 2734-2741. doi: 10.1016/j.jacc.2014.02.572
- 53- Mytas, D. Z., Stougiannos, P. N., Zairis, M. N., Foussas, S. G., Pyrgakis, V. N., & Kyriazis, I. A. (2009). Diabetic myocardial disease: pathophysiology, early diagnosis and therapeutic options. *J Diabetes Complications*, 23(4), 273-282. doi: 10.1016/j.jdiacomp.2007.12.005
- 54- Niture, S. K., Khatri, R., & Jaiswal, A. K. (2014). Regulation of Nrf2-an update. *Free Radic Biol Med*, 66, 36-44. doi: 10.1016/j.freeradbiomed.2013.02.008

- 55- Pal, P. B., Sonowal, H., Shukla, K., Srivastava, S. K., & Ramana, K. V. (2017). Aldose Reductase Mediates NLRP3 Inflammasome-Initiated Innate Immune Response in Hyperglycemia-Induced Thp1 Monocytes and Male Mice. *Endocrinology*, *158*(10), 3661-3675. doi: 10.1210/en.2017-00294
- 56- Passino, C., Barison, A., Vergaro, G., Gabutti, A., Borrelli, C., Emdin, M., & Clerico, A. (2015). Markers of fibrosis, inflammation, and remodeling pathways in heart failure. *Clin Chim Acta*, *443*, 29-38. doi: 10.1016/j.cca.2014.09.006
- 57- Pham, D., Albuquerque Rocha, N., McGuire, D. K., & Neeland, I. J. (2017). Impact of empagliflozin in patients with diabetes and heart failure. *Trends Cardiovasc Med*, *27*(2), 144-151. doi: 10.1016/j.tcm.2016.07.008
- 58- Pollack, P. S., Malhotra, A., Fein, F. S., & Scheuer, J. (1986). Effects of diabetes on cardiac contractile proteins in rabbits and reversal with insulin. *Am J Physiol*, *251*(2 Pt 2), H448-454. doi: 10.1152/ajpheart.1986.251.2.H448
- 59- Qi, Y., Xu, Z., Zhu, Q., Thomas, C., Kumar, R., Feng, H., . . . Guo, S. (2013). Myocardial loss of IRS1 and IRS2 causes heart failure and is controlled by p38alpha MAPK during insulin resistance. *Diabetes*, *62*(11), 3887-3900. doi: 10.2337/db13-0095
- 60- Qi, Y., Zhu, Q., Zhang, K., Thomas, C., Wu, Y., Kumar, R., . . . Guo, S. (2015). Activation of Foxo1 by insulin resistance promotes cardiac dysfunction and beta-myosin heavy chain gene expression. *Circ Heart Fail*, *8*(1), 198-208. doi: 10.1161/CIRCHEARTFAILURE.114.001457
- 61- Razeghi, P., Young, M. E., Cockrill, T. C., Frazier, O. H., & Taegtmeier, H. (2002). Downregulation of myocardial myocyte enhancer factor 2C and myocyte enhancer factor 2C-regulated gene expression in diabetic patients with nonischemic heart failure. *Circulation*, *106*(4), 407-411. doi: 10.1161/01.cir.0000026392.80723.d
- 62- Rosenkranz, A. C., Hood, S. G., Woods, R. L., Dusting, G. J., & Ritchie, R. H. (2003). B-type natriuretic peptide prevents acute hypertrophic responses in the diabetic rat heart: importance of cyclic GMP. *Diabetes*, *52*(9), 2389-2395. doi: 10.2337/diabetes.52.9.2389
- 63- Ryden, L., Armstrong, P. W., Cleland, J. G., Horowitz, J. D., Massie, B. M., Packer, M., & Poole-Wilson, P. A. (2000). Efficacy and safety of high-dose lisinopril in chronic heart failure patients at high cardiovascular risk, including those with diabetes mellitus. Results from the ATLAS trial. *Eur Heart J*, *21*(23), 1967-1978. doi: 10.1053/euhj.2000.2311
- 64- Sabolic, I., Vrhovac, I., Eror, D. B., Gerasimova, M., Rose, M., Breljak, D., . . . Koepsell, H. (2012). Expression of Na⁺-D-glucose

- cotransporter SGLT2 in rodents is kidney-specific and exhibits sex and species differences. *Am J Physiol Cell Physiol*, 302(8), C1174-1188. doi: 10.1152/ajpcell.00450.2011
- 65- Samuel, V. T., & Shulman, G. I. (2016). The pathogenesis of insulin resistance: integrating signaling pathways and substrate flux. *J Clin Invest*, 126(1), 12-22. doi: 10.1172/JCI77812
- 66- Sandesara, P. B., O'Neal, W. T., Kelli, H. M., Samman-Tahhan, A., Hammadah, M., Quyyumi, A. A., & Sperling, L. S. (2018). The Prognostic Significance of Diabetes and Microvascular Complications in Patients With Heart Failure With Preserved Ejection Fraction. *Diabetes Care*, 41(1), 150-155. doi: 10.2337/dc17-0755
- 67- Shome, J. S., Perera, D., Plein, S., & Chiribiri, A. (2017). Current perspectives in coronary microvascular dysfunction. *Microcirculation*, 24(1). doi: 10.1111/micc.12340
- 68- Song, R., Peng, W., Zhang, Y., Lv, F., Wu, H. K., Guo, J., . . . Xiao, R. P. (2013). Central role of E3 ubiquitin ligase MG53 in insulin resistance and metabolic disorders. *Nature*, 494(7437), 375-379. doi: 10.1038/nature11834
- 69- Strniskova, M., Barancik, M., Neckar, J., & Ravingerova, T. (2003). Mitogen-activated protein kinases in the acute diabetic myocardium. *Mol Cell Biochem*, 249(1-2), 59-65.
- 70- Sundgren, N. C., Giraud, G. D., Schultz, J. M., Lasarev, M. R., Stork, P. J., & Thornburg, K. L. (2003). Extracellular signal-regulated kinase and phosphoinositol-3 kinase mediate IGF-1 induced proliferation of fetal sheep cardiomyocytes. *Am J Physiol Regul Integr Comp Physiol*, 285(6), R1481-1489. doi: 10.1152/ajpregu.00232.2003
- 71- Talukder, M. A., Kalyanasundaram, A., Zuo, L., Velayutham, M., Nishijima, Y., Periasamy, M., & Zweier, J. L. (2008). Is reduced SERCA2a expression detrimental or beneficial to postischemic cardiac function and injury? Evidence from heterozygous SERCA2a knockout mice. *Am J Physiol Heart Circ Physiol*, 294(3), H1426-1434. doi: 10.1152/ajpheart.01016.2007
- 72- Tan, Y., Ichikawa, T., Li, J., Si, Q., Yang, H., Chen, X., . . . Cui, T. (2011). Diabetic downregulation of Nrf2 activity via ERK contributes to oxidative stress-induced insulin resistance in cardiac cells in vitro and in vivo. *Diabetes*, 60(2), 625-633. doi: 10.2337/db10-1164
- 73- Tate, M., Deo, M., Cao, A. H., Hood, S. G., Huynh, K., Kiriazis, H., . . . Ritchie, R. H. (2017). Insulin replacement limits progression of diabetic cardiomyopathy in the low-dose streptozotocin-induced diabetic rat. *Diab Vasc Dis Res*, 14(5), 423-433. doi: 10.1177/1479164117710390
- 74- Teshima, Y., Takahashi, N., Nishio, S., Saito, S., Kondo, H., Fukui, A., . . .

- . Saikawa, T. (2014). Production of reactive oxygen species in the diabetic heart. Roles of mitochondria and NADPH oxidase. *Circ J*, 78(2), 300-306. doi: 10.1253/circj.cj-13-1187
- 75- Thrainsdottir, I. S., Aspelund, T., Thorgeirsson, G., Gudnason, V., Hardarson, T., Malmberg, K., . . . Ryden, L. (2005). The association between glucose abnormalities and heart failure in the population-based Reykjavik study. *Diabetes Care*, 28(3), 612-616. doi: 10.2337/diacare.28.3.612
- 76- Uchimura, K., Hayata, M., Mizumoto, T., Miyasato, Y., Kakizoe, Y., Morinaga, J., . . . Kitamura, K. (2014). The serine protease prostaticin regulates hepatic insulin sensitivity by modulating TLR4 signalling. *Nat Commun*, 5, 3428. doi: 10.1038/ncomms4428
- 77- Vincent, M. A., Clerk, L. H., Lindner, J. R., Klibanov, A. L., Clark, M. G., Rattigan, S., & Barrett, E. J. (2004). Microvascular recruitment is an early insulin effect that regulates skeletal muscle glucose uptake in vivo. *Diabetes*, 53(6), 1418-1423. doi: 10.2337/diabetes.53.6.1418
- 78- Vincent, M. A., Clerk, L. H., Lindner, J. R., Price, W. J., Jahn, L. A., Leong-Poi, H., & Barrett, E. J. (2006). Mixed meal and light exercise each recruit muscle capillaries in healthy humans. *Am J Physiol Endocrinol Metab*, 290(6), E1191-1197. doi: 10.1152/ajpendo.00497.2005
- 79- Vollus, G. C., Bradley, E. A., Roberts, M. K., Newman, J. M., Richards, S. M., Rattigan, S., . . . Clark, M. G. (2007). Graded occlusion of perfused rat muscle vasculature decreases insulin action. *Clin Sci (Lond)*, 112(8), 457-466. doi: 10.1042/CS20060311
- 80- von Lewinski, D., Rainer, P. P., Gasser, R., Huber, M. S., Khafaga, M., Wilhelm, B., . . . Pieske, B. (2010). Glucose-transporter-mediated positive inotropic effects in human myocardium of diabetic and nondiabetic patients. *Metabolism*, 59(7), 1020-1028. doi: 10.1016/j.metabol.2009.10.025
- 81- Wakasaki, H., Koya, D., Schoen, F. J., Jirousek, M. R., Ways, D. K., Hoit, B. D., . . . King, G. L. (1997). Targeted overexpression of protein kinase C beta2 isoform in myocardium causes cardiomyopathy. *Proc Natl Acad Sci U S A*, 94(17), 9320-9325. doi: 10.1073/pnas.94.17.9320
- 82- Wang, J., Song, Y., Wang, Q., Kralik, P. M., & Epstein, P. N. (2006). Causes and characteristics of diabetic cardiomyopathy. *Rev Diabet Stud*, 3(3), 108-117. doi: 10.1900/RDS.2006.3.108
- 83- Wang, Y., Zhou, S., Sun, W., McClung, K., Pan, Y., Liang, G., . . . Cai, L. (2014). Inhibition of JNK by novel curcumin analog C66 prevents diabetic cardiomyopathy with a preservation of cardiac metallothionein expression. *Am J Physiol Endocrinol Metab*, 306(11),

- E1239-1247. doi:
10.1152/ajpendo.00629.2013
- 84- Westermeier, F., Riquelme, J. A., Pavez, M., Garrido, V., Diaz, A., Verdejo, H. E., . . . Lavandero, S. (2016). New Molecular Insights of Insulin in Diabetic Cardiomyopathy. *Front Physiol*, 7, 125. doi: 10.3389/fphys.2016.00125
- 85- Widyantoro, B., Emoto, N., Nakayama, K., Anggrahini, D. W., Adiarto, S., Iwasa, N., . . . Hirata, K. (2010). Endothelial cell-derived endothelin-1 promotes cardiac fibrosis in diabetic hearts through stimulation of endothelial-to-mesenchymal transition. *Circulation*, 121(22), 2407-2418. doi: 10.1161/CIRCULATIONAHA.110.938217
- 86- Xie, Z., Lau, K., Eby, B., Lozano, P., He, C., Pennington, B., . . . Zou, M. H. (2011). Improvement of cardiac functions by chronic metformin treatment is associated with enhanced cardiac autophagy in diabetic OVE26 mice. *Diabetes*, 60(6), 1770-1778. doi: 10.2337/db10-0351
- 87- Xu, Y. Z., Zhang, X., Wang, L., Zhang, F., Qiu, Q., Liu, M. L., . . . Wu, X. L. (2013). An increased circulating angiotensin II concentration is associated with hypoadiponectinemia and postprandial hyperglycemia in men with nonalcoholic fatty liver disease. *Intern Med*, 52(8), 855-861. doi: 10.2169/internalmedicine.52.8839
- 88- Yang, J., Park, Y., Zhang, H., Xu, X., Laine, G. A., Dellsperger, K. C., & Zhang, C. (2009). Feed-forward signaling of TNF-alpha and NF-kappaB via IKK-beta pathway contributes to insulin resistance and coronary arteriolar dysfunction in type 2 diabetic mice. *Am J Physiol Heart Circ Physiol*, 296(6), H1850-1858. doi: 10.1152/ajpheart.01199.2008
- 89- Yang, L., Zhao, D., Ren, J., & Yang, J. (2015). Endoplasmic reticulum stress and protein quality control in diabetic cardiomyopathy. *Biochim Biophys Acta*, 1852(2), 209-218. doi: 10.1016/j.bbadis.2014.05.006
- 90- Ye, G., Metreveli, N. S., Donthi, R. V., Xia, S., Xu, M., Carlson, E. C., & Epstein, P. N. (2004). Catalase protects cardiomyocyte function in models of type 1 and type 2 diabetes. *Diabetes*, 53(5), 1336-1343. doi: 10.2337/diabetes.53.5.1336
- 91- Young, M. E., Patil, S., Ying, J., Depre, C., Ahuja, H. S., Shipley, G. L., . . . Taegtmeier, H. (2001). Uncoupling protein 3 transcription is regulated by peroxisome proliferator-activated receptor (alpha) in the adult rodent heart. *FASEB J*, 15(3), 833-845. doi: 10.1096/fj.00-0351com
- 92- Zhang, H., Shi, X., Hampong, M., Blanis, L., & Pelech, S. (2001). Stress-induced inhibition of ERK1 and ERK2 by direct interaction with p38 MAP kinase. *J Biol Chem*, 276(10), 6905-6908. doi: 10.1074/jbc.C000917200
- 93- Zhang, Z., Wang, S., Zhou, S., Yan, X., Wang, Y., Chen, J., . . . Cai, L. (2014). Sulforaphane prevents the

- development of cardiomyopathy in type 2 diabetic mice probably by reversing oxidative stress-induced inhibition of LKB1/AMPK pathway. *J Mol Cell Cardiol*, 77, 42-52. doi: 10.1016/j.yjmcc.2014.09.022
- 94- Zhou, Y. T., Grayburn, P., Karim, A., Shimabukuro, M., Higa, M., Baetens, D., . . . Unger, R. H. (2000). Lipotoxic heart disease in obese rats: implications for human obesity. *Proc Natl Acad Sci U S A*, 97(4), 1784-1789. doi: 10.1073/pnas.97.4.1784
- 95- Zinman, B., Wanner, C., Lachin, J. M., Fitchett, D., Bluhmki, E., Hantel, S., . . . Investigators, E.-R. O. (2015). Empagliflozin, Cardiovascular Outcomes, and Mortality in Type 2 Diabetes. *N Engl J Med*, 373(22), 2117-2128. doi: 10.1056/NEJMoa1504720
- 96- Zou, M. H., & Xie, Z. (2013). Regulation of interplay between autophagy and apoptosis in the diabetic heart: new role of AMPK. *Autophagy*, 9(4), 624-625. doi: 10.4161/auto.23577